

Reduce your risk of chemotherapy-related infection with 1 of 3 administration options

The image is divided into three vertical panels. The left panel, titled "AT HOME", shows a woman lying on a couch using a tablet, with a Udenyca injection vial floating above her. The middle panel, titled "ON THE GO", shows a man talking on a phone while sitting at a desk with a laptop, with a Udenyca patch floating above him. The right panel, titled "IN THE OFFICE", shows a doctor in scrubs talking to two patients, with a Udenyca syringe floating above them. The text "IMMUNE SYSTEM SUPPORT THAT" is on the left, and "FITS INTO YOUR LIFE" is written in large, stylized letters across the bottom of the panels.

AT HOME

ON THE GO

IN THE OFFICE

IMMUNE SYSTEM SUPPORT THAT

FITS INTO YOUR LIFE

UDENYCA OFFERS ADMINISTRATION OPTIONS THAT MEET YOUR NEEDS

What is UDENYCA®?

UDENYCA is a prescription medicine used to help reduce the chance of infection due to a low white blood cell count, in people with certain types of cancer (non-myeloid), who receive anti-cancer medicines (chemotherapy) that can cause fever and low blood cell count.

IMPORTANT SAFETY INFORMATION

Do not take UDENYCA if you have had a serious allergic reaction to pegfilgrastim or filgrastim products.

Please see additional Important Safety Information throughout this brochure and click for the [Patient Information](#) and [Prescribing Information](#).


UDENYCA may help support your immune system while undergoing chemotherapy

CHEMOTHERAPY CAN DECREASE YOUR WHITE BLOOD CELL COUNT

Chemotherapy can help stop cancer cells from growing, but it can also lower your white blood cell count. A low white blood cell count can put you at risk for infection and a complication called febrile neutropenia, which is serious and can delay your chemotherapy treatments.

UDENYCA BOOSTS THE GROWTH OF YOUR WHITE BLOOD CELLS

UDENYCA increases your white blood cell count, which may help your body fight off infection and prevent febrile neutropenia.



THE MOST COMMON SIDE EFFECTS OF UDENYCA



BONE PAIN



ARM PAIN



LEG PAIN

These are not all of the possible side effects of UDENYCA. Please see the Important Safety Information throughout this brochure and contact your healthcare team for medical advice about side effects or concerns that you may have about your treatment.

Together, you can decide on the best way to manage side effects so you can stay on track with your treatment.

Please see the Important Safety Information throughout this brochure.

How will I know if I have an infection?

IT'S IMPORTANT TO TAKE YOUR TEMPERATURE

A fever is a sign of infection. Call your healthcare team immediately if you experience a fever of or higher than 100.4 °F (38.0 °C). Take your temperature any time you don't feel well or as recommended by your healthcare team.

Other signs of infection include chills, shortness of breath, sore throat, stiff neck, and burning or pain with urination. Talk to your healthcare team about signs to look out for.

If you develop a fever, it is a medical emergency, and you must call your healthcare team immediately or go to the emergency room at your local hospital.



Scan or click below to
VISIT [UDENYCA.COM](https://www.udenyca.com)
for more information
about your treatment

IMPORTANT SAFETY INFORMATION

What are possible side effects of UDENYCA?

UDENYCA may cause serious side effects, including:

- **Spleen rupture:** Your spleen may become enlarged and can rupture. A ruptured spleen can cause death. Call your healthcare provider right away if you have pain in the left upper stomach area or your left shoulder.
- **A serious lung problem called Acute Respiratory Distress Syndrome (ARDS):** Call your healthcare provider or get emergency care right away if you have shortness of breath with or without a fever, trouble breathing, or a fast rate of breathing.

Please see additional Important Safety Information throughout this brochure and click for the [Patient Information](#) and [Prescribing Information](#).

 **UDENYCA[®]**
pegfilgrastim-cbqv





The UDENYCA prefilled autoinjector

Treatment from the comfort of your home*



Delivers medicine
in less than 10 seconds†

*May also be given by your provider in their office.

†Please refer to the [Instructions for Use](#) before administering UDENYCA.

IMPORTANT SAFETY INFORMATION

What are possible side effects of UDENYCA? (Continued)

- **Serious allergic reactions:** UDENYCA can cause serious allergic reactions. These reactions can cause a rash over your whole body, shortness of breath, wheezing, dizziness, swelling around your mouth or eyes, fast heart rate, and sweating. **If you have any of these symptoms, stop using UDENYCA and call your healthcare provider or get emergency medical help right away.**
- **Sickle cell crises:** You may have a serious sickle cell crisis if you have a sickle cell disorder and receive UDENYCA. Serious sickle cell crises have happened in people with sickle cell disorders receiving pegfilgrastim that has sometimes led to death. Call your healthcare provider right away if you have symptoms of sickle cell crisis such as pain or difficulty breathing.

Click [here](#) or scan the QR code to **WATCH A VIDEO** to learn how to administer your medicine with the autoinjector



Please see additional Important Safety Information throughout this brochure and click for the [Patient Information](#), [Prescribing Information](#), and [Instructions for Use](#).

A quick guide to the UDENYCA autoinjector

Your healthcare provider will show you how to prepare and inject the UDENYCA autoinjector. ***This quick guide does not replace the Instructions for Use. Please refer to it before administering UDENYCA.*** Talk to your healthcare provider or pharmacist if you have any questions.

Only adults can self-inject UDENYCA with the autoinjector.

BEFORE GETTING STARTED

Make sure that you have these supplies ready:

- 1 UDENYCA autoinjector
- 1 alcohol wipe
- 1 cotton ball or gauze pad
- 1 FDA-approved sharps disposal container.

Learn more at [FDA.gov/safesharpsdisposal](https://www.fda.gov/safesharpsdisposal)


Take the UDENYCA autoinjector out of the refrigerator and bring to room temperature 30 minutes before use.

- Do not warm the autoinjector in any other way. For example, do not warm in a microwave, in hot water, or in direct sunlight
- Dispose of autoinjectors that have been left out at room temperature (68-77 °F) for more than 48 hours

Inspect your autoinjector and do not use if:


- The medicine appears discolored or cloudy
- The medicine contains lumps, flakes, or particles
- It has been dropped, crushed, or damaged in any way
- It is expired

ADMINISTERING UDENYCA




CHOOSE

Choose an injection site and **clean** it with an alcohol wipe. Stay 2 inches away from the navel if injecting into the abdomen. Clean and dry your hands.




REMOVE

With the clear cap of the autoinjector facing up, **pull** the cap straight off.



PRESS

Place the gray needle guard against your skin at a **90° angle**. **Press the autoinjector firmly** against the skin and you will hear the **first “click.”**



HOLD

Keep holding the autoinjector firmly against your skin until the orange indicator stops moving and you hear the **second “click.”** **After you hear the second “click,” continue to hold the autoinjector in place and count to 5.**

Please see the Important Safety Information throughout this brochure.



The UDENYCA on-body injector


**On-the-go treatment
injected in just 5 minutes***



Black **FILL INDICATOR** shows status of UDENYCA delivery

HIGH-QUALITY, WELL-TOLERATED ADHESIVE for secure attachment on your body

STATUS LIGHT signals if the on-body injector is working properly



FLASHING GREEN:

The on-body injector is working properly. **Do not** remove the on-body injector if the status light is flashing green.



SOLID GREEN (OR OFF):

Signals dose delivery is complete. Check to see if fill indicator reads empty.



FLASHING RED:

On-body injector error. If you hear beeping at any time, check the status light. If it is flashing red, call your healthcare provider right away as you may need a replacement dose.

*Please refer to the [Instructions for Use](#) before administering UDENYCA.

IMPORTANT SAFETY INFORMATION

What are possible side effects of UDENYCA? (Continued)

- **Allergies to Acrylics (UDENYCA ONBODY™ only):** The on-body injector (OBI) for UDENYCA uses acrylic adhesive. If you have had reactions to acrylic adhesives, use of this product may result in a significant reaction.
- **Kidney injury (glomerulonephritis):** UDENYCA can cause kidney injury. Call your healthcare provider right away if you develop any of the following symptoms:

- Swelling of your face or ankles
- Blood in your urine or dark colored urine
- You urinate less than usual
- **Increased white blood cell count (leukocytosis):** Your healthcare provider will check your blood during treatment with UDENYCA.

Please see additional Important Safety Information throughout this brochure and click for the [Patient Information](#), [Prescribing Information](#), and [Instructions for Use](#).

A quick guide to the UDENYCA on-body injector



The day you receive chemo, your on-body injector will be applied by your healthcare provider.



The day after chemo, your dose of UDENYCA will automatically be delivered about 27 hours after application.

Your healthcare provider will walk you through what to expect while wearing the device. ***This quick guide can help you understand what to expect while wearing the on-body injector, but it does not replace the Instructions for Use. Please refer to it while wearing the on-body injector.***


Talk to your healthcare provider or pharmacist if you have any questions.

Click [here](#) or scan the QR code to **WATCH A VIDEO** to learn what to expect with the on-body injector



AFTER YOUR PROVIDER APPLIES THE UDENYCA ON-BODY INJECTOR


1



MONITOR YOUR ON-BODY INJECTOR

For the next 27 hours, occasionally check the status light for at least 10 seconds. **Confirm it's flashing green.** If applied on the back of your arm, a caregiver must be available to check.


2



OBSERVE THE INJECTION

Before your injection begins, the on-body injector needle will automatically insert into your skin. During the injection, the on-body injector will flash a fast, green light and you may hear the pump working. It will take just 5 minutes for your dose of UDENYCA to be delivered.


3



CONFIRM INJECTION IS COMPLETE

When the injection is complete, a long beep will sound, and the status light will turn solid green. Confirm your injection is complete by checking to see that the status light is **solid green** or has switched off.

4



ENSURE YOUR ON-BODY INJECTOR IS EMPTY

Do not grasp the on-body injector itself to try and pull it off of your body. Instead, grab the edge of the adhesive and slowly peel the device off. Check to see if your on-body injector is empty. You should see a black line next to the empty indicator.


If you hear beeping at any time, check the status light. If it is flashing red, call your healthcare provider right away. Please see the Important Safety Information throughout this brochure.



The UDENYCA prefilled syringe

Injection administered by your healthcare team in the office

Click [here](#) or scan the QR code to **DISCOVER MORE** about the UDENYCA prefilled syringe



NEEDLE GUARD covers the needle and keeps you safe from needlesticks

A **LATEX-FREE** needle cap may reduce risk in people with latex sensitivity

The UDENYCA prefilled syringe—trusted administration for over 300,000 patients*

*January 2019 through November 2023.

IMPORTANT SAFETY INFORMATION

What are possible side effects of UDENYCA? (Continued)

- **Decreased platelet count (thrombocytopenia):** Your healthcare provider will check your blood during treatment with UDENYCA. Tell your healthcare provider if you have unusual bleeding or bruising during treatment with UDENYCA. This could be a sign of decreased platelet counts, which may reduce the ability of your blood to clot.
- **Capillary Leak Syndrome:** UDENYCA can cause fluid to leak from blood vessels into your body's tissues. This condition is called "Capillary Leak Syndrome" (CLS). CLS can quickly cause you to have symptoms that may become life-threatening. Get emergency medical help right away if you develop any of the following symptoms:

- Swelling or puffiness and are urinating less than usual
- Trouble breathing
- Swelling of your stomach-area (abdomen) and feeling of fullness
- Dizziness or feeling faint
- A general feeling of tiredness

Please see additional Important Safety Information throughout this brochure and click for the [Patient Information](#), [Prescribing Information](#), and [Instructions for Use](#).

Financial support is available

UDENYCA Solutions™, part of the Coherus Solutions™ family of support services, offers support for patients who are prescribed UDENYCA.



Working together to ensure patient access



Co-Pay Savings Program

The Co-Pay Savings Program may cover out-of-pocket expenses associated with UDENYCA for eligible patients with commercial insurance†

Pay as little as \$0 per dose of UDENYCA with a maximum annual benefit of \$15,000 for drug costs per calendar year†



Independent Foundation Support

UDENYCA Solutions may be able to help you find financial support through charitable foundations, or you can choose to contact these independent assistance foundations directly

Ask your doctor's office to help you apply for UDENYCA financial support.



FIND MORE DETAILS

about support and eligibility by visiting **UDENYCASolutions.com**, calling **1-844-4UDENYCA** (1-844-483-3692), or scanning this code

†Patients covered under Medicare, Medicaid, or any federal or state-funded program are not eligible for the Coherus Solutions Co-Pay Savings Program. The Co-Pay Savings Program ONLY covers the cost of UDENYCA and does not cover costs associated with the office visit. Eligibility requirements, type of assistance, and application process vary. Other restrictions apply; see Terms & Conditions at UDENYCASolutions.com. The Co-Pay Savings Program is subject to change or discontinuation without notice. This is not health insurance.

Please see the Important Safety Information throughout this brochure.



You're not in this alone

UDENYCA provides resources and support to help keep you on track with your treatment.



WHAT TO EXPECT DURING ADMINISTRATION

Click [here](#) or scan the code to watch a video to learn more about how UDENYCA is administered with the on-body injector and autoinjector.



FINANCIAL SUPPORT

Financial support may be available throughout your treatment journey with UDENYCA Solutions, part of the Coherus Solutions family of support services.



IMPORTANT SAFETY INFORMATION

What are possible side effects of UDENYCA? (Continued)

- **Myelodysplastic syndrome and acute myeloid leukemia:** If you have breast cancer or lung cancer, when UDENYCA is used with chemotherapy and radiation therapy, or with radiation therapy alone, you may have an increased risk of developing a precancerous blood condition called myelodysplastic syndrome (MDS) or a blood cancer called acute myeloid leukemia (AML). Symptoms of MDS and AML may include tiredness, fever, and easy bruising or bleeding. Call your healthcare provider if you develop these symptoms during treatment with UDENYCA.
- **Inflammation of the aorta (aortitis):** Inflammation of the aorta (the large blood vessel that transports blood from the heart to the body) has been reported in patients who received pegfilgrastim. Symptoms may include fever, abdominal pain, feeling tired, and back pain. Call your healthcare provider if you experience these symptoms.
- **Bone Imaging (Nuclear Imaging):** The formation of certain blood cells may increase and can be associated with short term changes in bone imaging. Your doctor should consider this when interpreting bone imaging results.
- **Allergies to Acrylics (UDENYCA ONBODY only):** The on-body injector (OBI) for UDENYCA uses acrylic adhesive. If you have had reactions to acrylic adhesives, use of this product may result in a significant reaction.

The most common side effects of UDENYCA are pain in the bones, arms, and legs.

These are not all the possible side effects of UDENYCA. Call your healthcare provider for medical advice about side effects. To report SUSPECTED ADVERSE REACTIONS, contact Coherus BioSciences at 1-800-4-UDENYCA (1-800-483-3692) or notify the FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

Please read the **Patient Information** and **Prescribing Information** for UDENYCA.



Coherus Solutions, UDENYCA Solutions, and UDENYCA ONBODY are trademarks, and UDENYCA is a registered trademark of Coherus BioSciences, Inc.

©2024 Coherus BioSciences, Inc. All rights reserved. US-UDY-0088 02/24